Climate Change Projections for Individual Tree Species

Northeast Ohio _____



This region's forests will be affected by a changing climate and other stressors during this century. Researchers and managers created an assessment that describes the vulnerability of forests in the Central Appalachians region (Butler et al. 2015: doi.org/10.2737/NRS-GTR-146). This report includes information on the current landscape, observed climate

trends, and a range of projected future climates. It also describes many potential climate change impacts to forests and summarizes key vulnerabilities for major forest ecosystems. This handout summarizes data from the U.S. Forest Service's Climate Change Tree Atlas (doi:org/10.2737/Climate-Change-Tree-Atlas-v4). Two climate scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The Tree Atlas provides information to interpret tree species changes:

- SUITABLE HABITAT calculated based on 45 variables that explain where conditions exist for a species, including soils, landforms, and climate variables.
- ADAPTABILITY based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- CAPABILITY a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (inventory data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species. Capability rating may not appropriately reflect the outlook for species with emerging severe forest health issues, such as ash species affected by emerald ash borer. See the table to the right for ratings.
- MIGRATION POTENTIAL MODEL when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management.

CLIMATE CHANGE CAPABILITY TABLE.

Capability is a rating of the species' ability to cope or persist with climate change. Species are organized into poor, fair, good, and mixed capability ratings. Species with new suitable habitat or low model reliability are excluded from this table. See the Tree Species Projections table legend on the following page for more information on ratings.

NOTE: Capability rating may not reflect severe forest health issues (e.g., emerald ash borer).

	,,					
POOR CAPABILITY						
American basswood	Quaking aspen					
American beech	Red pine					
Balsam fir	Scarlet oak					
Bigtooth aspen	Sourwood					
Black ash	Sweet birch					
Bur oak	Virginia pine					
Eastern hemlock	White ash					
Eastern white pine	Yellow birch					
Flowering dogwood						
FAIR CAPABILITY						
Black cherry	Osage-orange					
GOOD CAPABILITY						
Black oak	Pignut hickory					
Blackgum	Red maple					
Chinkapin oak	Sugar maple					
Hackberry	Sweetgum					
Mockernut hickory	White oak					
Northern red oak	Yellow-poplar					
MIXED CAPABILITY						
American elm	Shingle oak					
Shagbark hickory	<u> </u>					

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors, and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

CREDIT: This handout summarizes the full model results for the Northeast Ohio region. Data provided by the USDA Forest Service (M.P. Peters, A.M. Prasad, S.N. Matthews, & L.R. Iverson) as part of the Climate Change Tree Atlas (doi.org/10.2737/Climate-Change-Tree-Atlas-v4). Models and variables are described in Iverson et al. 2019 and Peters et al. 2019 (available at <u>fs.usda.gov/nrs/atlas/products/pubs</u>). More information on vulnerability and adaptation in the region can be found at <u>forestadaptation.org/central-appalachians</u>.







Tree Species Projections Table

Information presented in the table is from the Climate Change Tree Atlas regional summaries, more details at fs.usda.gov/nrs/atlas/combined/resources/summaries.

ADAPTABILITY: Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** Species may perform better than modeled
- · MEDIUM
- LOW Species may perform worse than modeled

HABITAT CHANGE: Projected change in suitable habitat between current and potential future conditions.

- ▲ INCREASE Projected increase of >20% by 2100
- ▼ **DECREASE** Projected decrease of >20% by 2100
- NO CHANGE Projected change of <20% by 2100
- ★ NEW HABITAT Tree Atlas projects new habitat for species not currently present

ABUNDANCE: Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + ABUNDANT
- · COMMON
- RARE

CAPABILITY: An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class, adaptability, and abundance within this region. Capability may not reflect severe forest health issues.

- △ **GOOD** Increasing suitable habitat, medium or high adaptability, and common or abundant
- FAIR Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability
- ▼ POOR Decreasing suitable habitat, medium or low adaptability, and uncommon or rare

				CLIMATE E (RCP 4.5)		H CLIMATE GE (RCP 8.5)				LOW CLIMATE CHANGE (RCP 4.5)		HIGH CLIMATE CHANGE (RCP 8.5)	
SPECIES	ADAPT	ABUN	HABITAT CHANGE		HABITA CHANG	T E CAPABILITY	SPECIES	ADAPT	ABUN	HABITAT CHANGE	Capability	HABITAT CHANGE	
American basswood	•	•	_	∇	_	∇	Osage-orange	+	_	•	0	•	0
American beech	•		_	∇	_	∇	Pignut hickory		_	<u> </u>	Δ	<u> </u>	Δ
American elm	•	•	•	0	_	Δ	Pin oak*	_	•	_	∇	_	$\overline{\nabla}$
American hornbeam*	•	_	_	∇	_	∇	Post oak	+		*		*	
Balsam fir	_	_	_	∇	_	$\overline{\nabla}$	Quaking aspen	•	•	_	∇	_	∇
Bigtooth aspen	•		_	∇	_	∇	Red maple	+	+	_	Δ	_	Δ
Bitternut hickory*	+	•	_	Δ	_	Δ	Red pine	_	_	_	∇	_	$\overline{\nabla}$
Black ash	_	_	_	∇	_	∇	Sassafras*	•	•	<u> </u>	Δ	<u> </u>	Δ
Black cherry	_	+	_	0	_	0	Scarlet oak	•	_	•	∇	•	$\overline{\nabla}$
Black locust*	•	•	_	Δ	_	Δ	Serviceberry*	•	_	_	∇	_	$\overline{\nabla}$
Black oak	•	•	_	Δ	_	Δ	Shagbark hickory	•		_	Δ	•	0
Black walnut*	•	•	_	Δ	_	Δ	Shingle oak	•	_	_	0	•	$\overline{\nabla}$
Black willow*	_	•	_	∇	_	∇	Shortleaf pine	•		*		*	
Blackgum	+	•	•	Δ	_	Δ	Shumard oak*	+		*		*	
Boxelder*	+	_	_	Δ	A	Δ	Silver maple*	+		•	Δ	•	Δ
Bur oak	+	_	_	∇	_	∇	Slippery elm*	•	•	_	Δ	<u> </u>	Δ
Chinkapin oak	•	_		Δ		Δ	Sourwood	+	_	_	∇	_	∇
Common persimmon*	+		*		*		Southern red oak	+		*		*	
Cucumbertree*	•	_	_	∇	_	∇	Sugar maple	+	+	•	Δ	_	Δ
Eastern cottonwood*	•	•	_	∇	_	∇	Swamp white oak*	•	•	_	∇	_	∇
Eastern hemlock	_	_		∇		∇	Sweet birch	_	_	_	∇	_	∇
Eastern hophornbeam'	* +	_		Δ		Δ	Sweetgum	•	_		Δ		Δ
Eastern redcedar	•		*		*		Sycamore*	•	_		Δ		Δ
Eastern white pine	_	•	_	∇		∇	Virginia pine	•	_	•	∇	•	∇
Flowering dogwood	•	_	•	∇		∇	White ash	_	•		∇	•	∇
Gray birch*	•	_	•	∇		∇	White oak	+	•		Δ		Δ
Green ash*	•	•		Δ		Δ	Winged elm	•		*		*	
Hackberry	+	_		Δ		Δ	Yellow birch	•	_	_	∇	_	∇
Mockernut hickory	+	_		Δ		Δ	Yellow-poplar	+	•		Δ	•	Δ
Northern red oak	+	•		Δ	•	Δ							