## Climate Change Projections for Individual Tree Species

## Southern and Coastal New England .



This region's forests will be affected by a changing climate and other stressors during this century. Researchers and managers created an assessment that describes the vulnerability of forests in the New England region (Janowiak et al. 2018: <a href="doi.org/10.2737/nrs-gtr-173">doi.org/10.2737/nrs-gtr-173</a>). This report includes information on the current landscape, observed climate trends, and a range of projected

future climates. It also describes many potential climate change impacts to forests and summarizes key vulnerabilities for major forest ecosystems. This handout summarizes data from the U.S. Forest Service's Climate Change Tree Atlas (doi.org/10.2737/Climate-Change-Tree-Atlas-v4). Two climate scenarios are presented to "bracket" a range of possible futures. These future climate projections (2070 to 2099) provide information about how individual tree species may respond to a changing climate. Results for "low" and "high" emissions scenarios can be compared on the reverse side of this handout.

The Tree Atlas provides information to interpret tree species changes:

- SUITABLE HABITAT calculated based on 45 variables that explain where conditions exist for a species, including soils, landforms, and climate variables.
- ADAPTABILITY based on life-history traits that might increase or decrease tolerance of expected changes, such as the ability to withstand different forms of disturbance.
- CAPABILITY a rating of the species' ability to cope or persist with climate change in this region based on suitable habitat change (statistical modeling), adaptability (literature review and expert opinion), and abundance (inventory data). The capability rating is modified by abundance information; ratings are downgraded for rare species and upgraded for abundant species. Capability rating may not appropriately reflect the outlook for species with emerging severe forest health issues, such as ash species affected by emerald ash borer. See the table to the right for ratings.
- **MIGRATION POTENTIAL MODEL** when combined with habitat suitability, an estimate of a species' colonization likelihood for new habitats. This rating can be helpful for assisted migration or focused management.

Remember that models are just tools, and they're not perfect. Model projections can't account for all factors that influence future species success. If a species is rare or confined to a small area, model results may be less reliable. These factors,

## CLIMATE CHANGE CAPABILITY TABLE.

Capability is a rating of the species' ability to cope or persist with climate change. Species are organized into poor, fair, good, and mixed capability ratings. Species with new suitable habitat or low model reliability are excluded from this table. See the Tree Species Projections table legend on the following page for more information on ratings.

**NOTE:** Capability rating may not reflect severe forest health issues (e.g., emerald ash borer).

reater issues (e.g., emerate	,
POOR CAPABILITY	
American basswood	Northern white-ceda
Balsam fir	Paper birch
Balsam poplar	Pitch pine
Black ash	Red pine
Black spruce	Red spruce
Bur oak	Striped maple
Eastern white pine	Sweet birch
Hackberry	Tamarack (native)
Northern pin oak	White spruce
FAIR CAPABILITY	
Quaking aspen	White ash
GOOD CAPABILITY	
American beech	Pignut hickory
American holly	Post oak
Black cherry	Red maple
Black oak	Scarlet oak
Blackgum	Shagbark hickory
Chestnut oak	Sugar maple
Eastern redcedar	Sweetgum
Mockernut hickory	White oak
Northern red oak	Yellow-poplar
MIXED CAPABILITY	
American elm	Flowering dogwood
Bigtooth aspen	Yellow birch
Eastern hemlock	

and others, could cause a particular species to perform better or worse than a model projects. Human choices will also continue to influence forest distribution, especially for tree species that are projected to increase. Planting programs may assist the movement of future-adapted species, but this will depend on management decisions. Despite these limits, models provide useful information about future expectations. It's perhaps best to think of these projections as indicators of possibility and potential change.

**CREDIT:** This handout summarizes the full model results for the Southern and Coastal New England region. Data provided by the USDA Forest Service (M.P. Peters, A.M. Prasad, S.N. Matthews, & L.R. Iverson) as part of the Climate Change Tree Atlas (doi.org/10.2737/Climate-Change-Tree-Atlas-v4). Models and variables are described in Iverson et al. 2019 and Peters et al. 2019 (available at <a href="mailto:fs.usda.gov/nrs/atlas/products/pubs">fs.usda.gov/nrs/atlas/products/pubs</a>). More information on vulnerability and adaptation in the region can be found at <a href="mailto:forestadaptation.org/new-england">forestadaptation.org/new-england</a>.







## Tree Species Projections Table

Information presented in the table is from the Climate Change Tree Atlas regional summaries, more details at fs.usda.gov/nrs/atlas/combined/resources/summaries.

**ADAPTABILITY:** Life-history factors, such as the ability to respond favorably to disturbance, that are not included in the Tree Atlas model and may make a species more or less able to adapt to future stressors.

- + **HIGH** Species may perform better than modeled
- · MEDIUM
- LOW Species may perform worse than modeled

**HABITAT CHANGE:** Projected change in suitable habitat between current and potential future conditions.

- ▲ INCREASE Projected increase of >20% by 2100
  - **DECREASE** Projected decrease of >20% by 2100
- NO CHANGE Projected change of <20% by 2100
- ★ NEW HABITAT Tree Atlas projects new habitat for species not currently present

**ABUNDANCE:** Based on Forest Inventory Analysis (FIA) summed Importance Value data, calibrated to a standard geographic area.

- + ABUNDANT
- COMMON
- RARE

**CAPABILITY:** An overall rating that describes a species' ability to cope or persist with climate change based on suitable habitat change class, adaptability, and abundance within this region. Capability may not reflect severe forest health issues.

- △ **GOOD** Increasing suitable habitat, medium or high adaptability, and common or abundant
- FAIR Mixed combinations, such as a rare species with increasing suitable habitat and medium adaptability
- ▼ POOR Decreasing suitable habitat, medium or low adaptability, and uncommon or rare

							and uncommon o						
				LIMATE (RCP 4.5)		CLIMATE E (RCP 8.5)					CLIMATE E (RCP 4.5)		CLIMATE E (RCP 8.5)
SPECIES	ADAPT	ABUN	HABITAT CHANGE O	CAPABILITY	HABITAT CHANGE	Capability	SPECIES	ADAPT	ABUN	HABITAT CHANGE		HABITAT CHANGE	Capability
American basswood	•	_	•	$\nabla$	•	$\nabla$	Northern red oak	+	+	•	Δ	_	Δ
American beech	•	•	<u> </u>	Δ	<u> </u>	Δ	Northern white-cedar	•	_	_	$\nabla$	_	$\nabla$
American elm	•	•	_	$\nabla$	•	0	Paper birch	•	•	_	$\nabla$	_	$\nabla$
American holly	•	_	<u> </u>	Δ	<u> </u>	Δ	Pignut hickory	•	•	_	Δ		Δ
American hornbeam*	•	_	•	$\nabla$	_	Δ	Pin cherry*	•	_	_	$\nabla$	_	$\overline{\nabla}$
American mountain-ash	* –	_	_	$\nabla$	_	$\nabla$	Pin oak*	_	_	•	$\nabla$		0
Atlantic white-cedar*	_	_	•	$\nabla$	•	$\nabla$	Pitch pine	•	•	_	$\overline{\nabla}$	_	$\overline{\nabla}$
Bald cypress	•		*		*		Post oak	+	_	<u> </u>	Δ	_	Δ
Balsam fir	_	•	_	$\nabla$	_	$\nabla$	Quaking aspen	•	•	•	0	•	0
Balsam poplar	•	_	_	$\nabla$	_	$\nabla$	Red maple	+	+	_	Δ	_	Δ
Bigtooth aspen	•	•	_	Δ	•	0	Red pine	_	_	_	$\nabla$	_	$\overline{\nabla}$
Bitternut hickory*	+	_	•	0	_	Δ	Red spruce	_	_	_	$\nabla$	_	$\overline{\nabla}$
Black ash	_	_	_	$\nabla$	_	$\nabla$	Sassafras*		_	_	Δ	<u> </u>	Δ
Black cherry	_	•	_	Δ	_	Δ	Scarlet oak	•		_	Δ	_	Δ
Black locust*		_	_	0	_	0	Shagbark hickory	•		_	Δ	<u> </u>	Δ
Black oak	•	•	_	Δ	_	Δ	Shortleaf pine	•		*		*	
Black spruce		_	_	$\nabla$	_	$\nabla$	Silver maple*	+	_	_	$\nabla$	•	0
Black walnut*	•	_	_	$\nabla$	•	$\nabla$	Slippery elm*		_	_	$\nabla$	_	$\overline{\nabla}$
Blackgum	+	_	_	Δ	_	Δ	Southern red oak	+		*		*	
Boxelder*	+	_	•	0	•	0	Striped maple	•	_	_	$\nabla$	_	$\nabla$
Bur oak	+	_	_	$\nabla$	_	$\nabla$	Sugar maple	+		<u> </u>	Δ		Δ
Chestnut oak	+	_	_	Δ	_	Δ	Swamp chestnut oak*		_	_	$\nabla$	_	Δ
Chinkapin oak	•				*		Swamp tupelo	_		*		*	
Eastern hemlock	_	+	_	0	_	$\nabla$	Swamp white oak*	•	_	•	$\nabla$	•	$\overline{\nabla}$
Eastern redcedar	•	•	_	Δ	_	Δ	Sweet birch	-	•	•	$\nabla$	_	$\nabla$
Eastern white pine	_	+	_	$\nabla$	_	$\nabla$	Sweetbay	•		*		*	
Flowering dogwood		_	•	$\nabla$	<u> </u>	0	Sweetgum	•	_	<u> </u>	Δ	<u> </u>	Δ
Gray birch*	•	_	•	$\nabla$	<u> </u>	0	Sycamore*	•	_	•	$\nabla$	<u> </u>	0
Green ash*	•	_	<u> </u>	0	<u> </u>	Δ	Tamarack (native)	_	_	_	$\nabla$	_	$\nabla$
Hackberry	+	_	_	$\nabla$	_	$\overline{\nabla}$	Virginia pine	•		*		*	
Honeylocust*	+	_	_	$\nabla$	_	$\nabla$	Water hickory	•				*	
Ironwood*	+	_	•	0	_	Δ	White ash	_		_	0	_	0
Loblolly pine	•		*		*		White oak	+		_	Δ		Δ
Longleaf pine			*		*		White spruce	•	_	_	$\overline{\nabla}$	_	$\overline{\nabla}$
Mockernut hickory	+	_	<u> </u>	Δ	_	Δ	Yellow birch	•	•	•	0	<u> </u>	Δ
Northern pin oak	+	_		$\overline{\nabla}$	_	$\overline{\nabla}$	Yellow-poplar	+	_	_	Δ		
				•		•	P. P. P.				<del></del>		<del></del>